

Insulator-Metal Transition in the One and Two-Dimensional Hubbard Models

F.F. Assaad and M. Imada

Institute for Solid State Physics, University of Tokyo,
7-22-1 Roppongi, Minato-ku, Tokyo 106, Japan.

Abstract

We use Quantum Monte Carlo methods to determine $T = 0$ Green functions, $G(\vec{r}, \omega)$, on lattices up to 16×16 for the 2D Hubbard model at $U/t = 4$. For chemical potentials, μ , within the Hubbard gap, $|\mu| < \mu_c$, and at *long* distances, $\vec{r}$, $G(\vec{r}, \omega = \mu) \sim e^{-|\vec{r}|/\xi_l}$ with critical behavior: $\xi_l \sim |\mu - \mu_c|^{-\nu}$, $\nu = 0.26 \pm 0.05$. This result stands in agreement with the assumption of hyper-scaling with correlation exponent $\nu = 1/4$ and dynamical exponent $z = 4$. In contrast, the generic band insulator as well as the metal-insulator transition in the 1D Hubbard model are characterized by $\nu = 1/2$ and $z = 2$.

PACS numbers: 71.27.+a, 71.30.+h, 71.10.+x

At zero-temperature a continuous metal-insulator transition driven by a change in chemical potential may be characterized by the compressibility, χ_c , or the Drude weight, D . In the Mott insulating phase both D and χ_c vanish while they remain finite in the metallic phase [1,2]. In order to describe the metal-insulator transition from the insulator side, we consider the zero-temperature Green function $G(\vec{r}, \omega)$ [3]. At *long* distances, $|\vec{r}|$, and for values of the chemical potential, μ , within the charge gap, $G(\vec{r}, \omega = \mu) \sim e^{-|\vec{r}|/\xi_l}$. The metal-insulator transition may be characterized by the divergence of ξ_l as the critical chemical potential, μ_c , is approached from the insulating phase. ξ_l may be interpreted as the localization length involved in transferring a particle over a distance $\vec{r}$ from the electronic system to the heat bath lying at energy μ within the charge gap. Under the assumption of hyperscaling, the above quantities are expected to satisfy the scaling relations:

$$\xi_l \sim \Delta^{-\nu}, \quad \chi_c \sim \Delta^{-\nu(z-d)}, \quad D \sim \Delta^{\nu(d+z-2)}, \quad (1)$$

where $\Delta = |\mu - \mu_c|$, d is the dimensionality and ν (z) corresponds to the correlation length (dynamical) exponent [2]. Since the control parameter, Δ , corresponds to the chemical potential, one obtains the additional constraint $\nu z = 1$ as well as $\delta \sim \Delta^{\nu(d+z)-1}$, δ being the doping concentration. In this letter, based on a recently developed numerically stable Quantum Monte Carlo (QMC) algorithm to calculate zero-temperature imaginary time Green functions [4], we calculate ξ_l for the two-dimensional repulsive Hubbard model at $U/t = 4$. We obtain: $\mu_c = 0.67 \pm 0.015$ in units of the hopping matrix element and $\xi_l \sim |\mu - \mu_c|^{-\nu}$ with $\nu = 0.26 \pm 0.05$. On the other hand, the compressibility data of Furukawa and Imada [5] in the metallic phase is consistent with $\chi_c \sim |\mu - \mu_c|^{-1/2}$. When hyperscaling describes the transition, the above scaling of the compressibility also leads to $\nu = 1/4$ [2]. Comparison of those results puts the hyperscaling assumption on a firmer basis. The present estimate of ν is a more direct determination of the characteristic length scale. In contrast, the Mott transition in the one-dimensional Hubbard model satisfies the scaling relations (1) with $\nu = 1/2$ and $z = 2$ [2,6–8]. The generic band insulator in all dimensions equally belongs to the universality class $\nu = 1/2$ and $z = 2$ [2]. The above results point out

the anomalous character of the Mott transition in the two-dimensional Hubbard model.

The Hubbard model we consider reads:

$$H - \mu N = -t \sum_{\langle \vec{i}, \vec{j} \rangle, \sigma} c_{i,\sigma}^\dagger c_{j,\sigma} + U \sum_{\vec{i}} (n_{i,\uparrow} - \frac{1}{2})(n_{i,\downarrow} - \frac{1}{2}) - \mu \sum_{\vec{i}, \sigma} c_{i,\sigma}^\dagger c_{i,\sigma} \quad (2)$$

Here, $\langle \vec{i}, \vec{j} \rangle$ denotes nearest-neighbors. $c_{i,\sigma}^\dagger$ ($c_{i,\sigma}$) creates (annihilates) an electron with z -component of spin σ on site $\vec{i}$ and $n_{i,\sigma} = c_{i,\sigma}^\dagger c_{i,\sigma}$. In this notation half-band filling corresponds to $\mu = 0$. We start by considering the non-interacting case, $U/t = 0$. In Fourier space, the single particle energies are given by $\epsilon_{\vec{k}} = -2t(\cos(\vec{k}\vec{a}_x) + \cos(\vec{k}\vec{a}_y))$, $\vec{a}_x, \vec{a}_y$ being the lattice constants. In this letter, the length scale is set by: $|\vec{a}_x| = |\vec{a}_y| = 1$. At zero temperature, an insulator-metal transition will occur when $\mu \rightarrow \mu_c = 4t$. For those chemical potentials, the zero-temperature Green function [3] at $\omega = \mu$ is given by:

$$G(\vec{r}, \omega = \mu) = \frac{2}{N} \sum_{\vec{k}} \frac{e^{i\vec{k}\vec{r}}}{\epsilon_{\vec{k}} - \mu} \quad (3)$$

where N denotes the number of sites of the square lattice and the factor 2 corresponds to the summation over the spin degrees of freedom. Numerically, one obtains: $G(\vec{r}, \omega = \mu) \sim e^{-|\vec{r}|/\xi_l}$ with critical behavior: $\xi_l \sim |\mu - \mu_c|^{-1/2}$ [9]. One may check that this example of a band insulator-metal transition satisfies the above scaling relations (1) and is characterized by the exponent $\nu = 1/2$. At finite values of U/t and half-band filling the antiferromagnetic Hartree-Fock approximation equally yields the exponent $\nu = 1/2$ for ξ_l . However, this approximation does not satisfy the hyperscaling assumption.

The physical interpretation of ξ_l is facilitated by considering the single impurity Hamiltonian (Fano-Anderson model) [10,11]:

$$H = \sum_{\vec{k}} \epsilon_{\vec{k}} c_{\vec{k}}^\dagger c_{\vec{k}} + \epsilon_b b^\dagger b + \frac{t_b}{\sqrt{N}} \sum_{\vec{k}} (c_{\vec{k}}^\dagger b + b^\dagger c_{\vec{k}}). \quad (4)$$

Here, $b^\dagger$ creates an electron in the impurity state at the origin and energy ϵ_b . The hybridization between the localized state and the band electrons alters the energy of the impurity level to the value: $E_b = \epsilon_b + \frac{t_b^2}{N} \sum_{\vec{k}} \frac{1}{E_b - \epsilon_{\vec{k}}}$. We will assume $E_b > \epsilon_{\vec{k}}$ for all $\vec{k}$. When all single

particle states of the valence band are filled and the impurity single particle state empty, the probability amplitude of transferring a band electron at site $\vec{r}$ to the impurity state is:

$$\frac{\langle \Psi_0 | b^\dagger c_{\vec{r}} | \Psi_0 \rangle}{\langle \Psi_0 | \Psi_0 \rangle} = -\alpha(E_b) \frac{t_b}{N} \sum_{\vec{k}} \frac{e^{i\vec{k}\vec{r}}}{\epsilon_{\vec{k}} - E_b} \quad (5)$$

where the normalization factor is given by $\alpha^{-1}(E_b) = 1 + \frac{t_b^2}{N} \sum_{\vec{k}} \frac{1}{(E_b - \epsilon_{\vec{k}})^2}$. Comparison between equations (5) and (3) show that the spatial dependence of the two quantities is identical. ξ_l may thus be interpreted as the localization length involved in transferring a particle over a distance $\vec{r}$ from the valence band to the heat bath (see equation (3)), or to the impurity state (see equation (5)). This definition of ξ_l bears some similarity to that applied in finite size scaling studies of Anderson localized states [12,13]. When the imaginary part of the Green function vanishes in the insulating phase, ξ_l may be used to study the metal-insulator transition.

To obtain an estimate of the critical exponent ν for the Hubbard model, we require an accurate determination of the critical chemical potential, μ_c as well as of the localization length, ξ_l . Both quantities may be obtained from the knowledge of:

$$G_\sigma(\vec{r}, \tau) = \Theta(\tau) \frac{\langle \Psi_0 | c_{\vec{r},\sigma}(\tau) c_{\vec{0},\sigma}^\dagger | \Psi_0 \rangle}{\langle \Psi_0 | \Psi_0 \rangle} - \Theta(-\tau) \frac{\langle \Psi_0 | c_{-\vec{r},\sigma}^\dagger(-\tau) c_{\vec{0},\sigma} | \Psi_0 \rangle}{\langle \Psi_0 | \Psi_0 \rangle}, \quad (6)$$

where $c_{\vec{r},\sigma}(\tau) = e^{\tau H} c_{\vec{r},\sigma} e^{-\tau H}$. Here $|\Psi_0\rangle$ denotes the ground state of the half-filled ($\mu = 0$) Hubbard Hamiltonian (2). The above quantity may be efficiently calculated with QMC methods. Since we are at half-band filling, the sign problem does not occur and we are able to consider lattice sizes up to linear dimension $L = 16$, namely $N = 16 \times 16$ without any serious difficulties. The calculation of imaginary time Green function in the zero-temperature auxiliary field QMC algorithm [14,15] was first reported by Deisz et al. [16]. However, their approach does not incorporate a numerical stabilization scheme and they are thus restricted to small values of τ (i.e. $\tau t \sim 2.5$). The authors have developed a numerically stable QMC algorithm for the calculation of $G_\sigma(\vec{r}, \tau)$. The details of the algorithm may be found in reference [4]. All our calculations were performed with periodic boundary conditions.

From the knowledge of $G_\sigma(\vec{r}, \tau)$ on an N -site lattice, we may obtain an estimate of the charge gap. We denote by $|\Psi_n^N\rangle$ the eigenvector of the Hamiltonian H with eigenvalue E_n^N in the N -particle Hilbert space. With this notation,

$$G(\vec{r}=0, \tau) \equiv \sum_\sigma G_\sigma(\vec{r}=0, \tau) = \frac{1}{N} \sum_{n, \vec{i}, \sigma} |\langle \Psi_0^N | c_{\vec{i}, \sigma} | \Psi_n^{N+1} \rangle|^2 \exp\left(-\tau (E_n^{N+1} - E_0^N)\right). \quad (7)$$

for $\tau > 0$. Fig. 1a, plots $G(\vec{r}=0, \tau)$ for a 16×16 lattice at $U/t = 4$. We may obtain a reliable estimate of the charge gap for this lattice size by fitting $G(\vec{r}=0, \tau)$ to the form $e^{-\Delta_c \tau}$ with $\Delta_c \equiv E_0^{N+1} - E_0^N$ for *large* values of τ . Fig. 1b shows Δ_c as a function of linear lattice size. Extrapolation to the thermodynamic limit yields: $\Delta_c/t = 0.67 \pm 0.015$. This result stands in good agreement with the quoted result of Furukawa and Imada [5]: $\Delta_c/t = 0.58 \pm 0.08$. Since in the notation of equation (2) the Hubbard model satisfies particle-hole symmetry at $\mu = 0$, the critical chemical potential is nothing but the charge gap: $\mu_c \equiv \Delta_c$.

For values of the chemical potential within the charge gap, $|\mu| < \mu_c$, the zero temperature Green function is real and may be obtained through the relation:

$$G(\vec{r}, \omega = \mu) = - \int_{-\infty}^{\infty} d\tau G(\vec{r}, \tau) e^{\tau \mu}. \quad (8)$$

The Green function $G(\vec{r}, \tau)$ is computed at half-band filling where the sign problem is not present and the statistical uncertainty does not grow exponentially with lattice size. However, since we are multiplying the QMC data by the factor $e^{\tau \mu}$, the statistical uncertainty will grow exponentially with increasing values of τ for $\tau \mu > 0$. For each lattice size L , we have considered the largest distance $\vec{r} = (L/2, L/2)$. For this distance, $G(\vec{r} = (L/2, L/2), \tau)$ is plotted in Fig. 2. Due to particle-hole symmetry at $\mu = 0$, $G(\vec{r} = (L/2, L/2), \tau) = -G(\vec{r} = (L/2, L/2), -\tau)$. For the imaginary time integration, (see equation (8)) and values of the chemical potential $|\mu| < 0.65t$, a cutoff of $\tau t = 10$ proved to be sufficient for the determination of the Green function (see Fig. 2). $G(\vec{r} = (L/2, L/2), \omega = \mu)$ as a function of lattice size is plotted in Fig. 3 for several values of μ . For lattice sizes ranging from $L = 6$ to $L = 16$, an exponential decay may be seen within the quoted statistical uncertainty. From this data, we obtain an estimate of the localization length ξ_L . With the above determined

value of μ_c , we plot $|\mu - \mu_c|$ versus ξ_l (see Fig. 4) on a log-log plot. For all considered chemical potentials, $\xi_l/a < 8\sqrt{2}$ which corresponds to our largest considered distance ($L = 16$). The slopes in Fig. 4 correspond to values of the critical exponent $\nu = 1/2$ and $\nu = 1/4$. The QMC data is consistent with $\nu = 1/4$ and seems to rule out the possibility $\nu = 1/2$. A statistical analysis yields: $\nu = 0.26 \pm 0.05$.

The choice $\vec{r} = (L/2, L/2)$ is very convenient since apart from a sign change between $L = 4n + 2$ and $L = 4n$ lattices the exponential decay of $G(\vec{r} = (L/2, L/2), \omega = \mu)$ as a function of L , is not masked by a non-trivial oscillation. For other choices of $\vec{r}$ we expect to obtain the same results since due to the point group symmetry of the square lattice, all $\vec{k}$ -points in the Brillouin zone contribute to $G(\vec{r} = (L/2, L/2), \tau)$. We could however not check this point numerically.

To confirm the validity of our approach, we consider the Mott transition in the one-dimensional Hubbard model at $U/t = 4$. A similar QMC calculation as described above for the two-dimensional case yields a value of the correlation length exponent consistent with $\nu = 1/2$ (see Fig. 5). We have obtained $\mu_c/t = 0.66 \pm 0.015$ which is consistent with the exact result of Lieb and Wu [17]: $\mu_c/t = 0.643$. Chains of linear length up to $L = 24$ were considered. This result stands in agreement with the assumption of hyperscaling with exponents $\nu = 1/2$ and $z = 2$ [2,6-8].

In conclusion, we have determined the correlation length exponent from the knowledge of ξ_l in the insulating phase of the two-dimensional Hubbard model and obtained: $\nu = 0.26 \pm 0.05$. Under the assumption of hyperscaling with exponents $\nu = 1/4$ and $z = 4$, this result stands in agreement with compressibility measurements in the metallic phase [5]. We have shown that a similar calculation for the one-dimensional Hubbard model, yields results consistent with $\nu = 1/2$. Several anomalous aspects of the metal-insulator transition are inferred when it is characterized by the universality class, $\nu = 1/4$ and $z = 4$ [2]. Based on a single-particle theory, the exponents $\nu = 1/4$, $z = 4$ are consistent with the interpretation of the Mott transition driven by the divergence of the effective mass as opposed to the vanishing of the number of charge carriers. This statement is supported by

the compressibility in the metallic phase [5] as well as by the high frequency Hall coefficient [18]. Another consequence of the exponent $\nu = 1/4$, is the behavior of the Drude weight in the vicinity of the Mott transition: $D \sim \delta^2$, δ being the hole-density. As a by-product, we have produced an accurate estimate of the charge gap for the two-dimensional Hubbard model at $U/t = 4$: $\Delta_c/t = 0.67 \pm 0.015$. From the technical point of view, we have introduced an efficient method to obtain information on the nature of the metal-insulator transition by approaching the transition from the insulator side. The most important fact, is that for models which show particle-hole symmetry, such as dimerized Hubbard models, the here presented method is not plagued by the sign problem and large lattice sizes may be considered.

F.F.A. thanks the JSPS for financial support. The numerical calculations were carried out on the Fujitsu VPP500 of the Supercomputer Center of the Institute for Solid State Physics, Univ. of Tokyo. This work is supported by a Grant-in-Aid for Scientific Research on the Priority Area ‘Anomalous Metallic State near the Mott Transition’ from the Ministry of Education, Science and Culture, Japan.

REFERENCES

- [1] W. Kohn, Phys. Rev. **133A**, 171, (1964).
- [2] M. Imada, J. Phys. Soc. of Jpn. **64**, 2954 (1995).
- [3] We define the zero-temperature Green function by: $iG(\vec{r}, \omega) = \sum_{\sigma} \int dt e^{i\omega t} \langle \Psi_0 | T c_{\sigma, \vec{r}}(t) c_{\sigma, \vec{0}}^{\dagger}(0) | \Psi_0 \rangle$. Here, the notation is standard [11] and $|\Psi_0\rangle$ corresponds to the ground state in the insulating phase.
- [4] F.F. Assaad and M. Imada, to appear in J. Phys. Soc. Jpn. (cond-mat/9508113)
- [5] N. Furukawa and M. Imada, J. Phys. Soc. Jpn. **62**, 2557, (1993).
- [6] M. Mori, H. Fukuyama and M. Imada, J. Phys. Soc. of Jpn. **63**, 1639, (1994).
- [7] B.S. Shastry and B. Sutherland, Phys. Rev. Lett. **65**, 243, (1990).
- [8] M. Imada, J. Phys. Soc. Jpn. **63**, 3059, (1994).
- [9] For the dispersion relation $\epsilon_{\vec{k}} = \frac{\vec{k}^2}{2m}$ the integration may be done analytically to obtain the result $\xi_l \sim |\mu - \mu_c|^{-1/2}$, with $\mu_c = 0$.
- [10] P.W. Anderson, Phys. Rev. **124**, 41, (1961).
- [11] G.D. Mahan, Many-Particle Physics, Plenum Press, New York, (1981).
- [12] D.C. Herbert and R. Jones, J. Phys. C **4**, 1145, (1971).
- [13] A. MacKinnon and B. Kramer, Phys. Rev. Lett. **21**, 1546, (1981).
- [14] S. Sorella, S. Baroni, R. Car, and M. Parrinello, Europhys. Lett. **8** (1989) 663. S. Sorella, E. Tosatti, S. Baroni, R. Car, and M. Parrinello, Int. J. Mod. Phys. B **1** (1989) 993.
- [15] G. Sugiyama and S.E. Koonin, Anals of Phys. **168** (1986) 1.
- [16] J.J. Deisz, W. von der Linden, R. Preuss and W. Hanke, to appear in *Computer simulations in Condensed Matter Physics VIII*, Eds. D.P. Landau, K.K. Mon and H.B.

Schüttler (Springer Verlag, Heidelberg, Berlin, 1995).

[17] E.H. Lieb and F.Y. Wu, Phys. Rev. Lett. **20**, 1445, (1968).

[18] F.F. Assaad and M. Imada, Phys. Rev. Lett. **74**, 3872, (1995).

Figure captions

Fig. 1 a) $\ln G(\vec{r} = 0, \tau)$ as a function of τ for the half-filled ($\mu = 0$) 2D Hubbard model at $U/t = 4$ on a 16×16 lattice. The solid line corresponds to a least square fit of $G(\vec{r} = 0, \tau)$ to the form $\exp(-\Delta_c \tau)$ at *large* values of τ . b) Δ_c as a function of linear lattice size L . The solid circle at $1/L = 0$ corresponds to Δ_c as obtained in reference [5].

Fig. 2 $|G(\vec{r} = (L/2, L/2), \tau)|$ as a function of system size and imaginary time τ for the 2D Hubbard model.

Fig. 3 $\ln |G(\vec{r} = (L/2, L/2), \omega = \mu)|$ as a function of distance and chemical potential for the 2D Hubbard model. The solid lines correspond to a least square fit of $|G(\vec{r} = (L/2, L/2), \omega = \mu)|$ to the form $\exp(-|\vec{r}|/\xi_l)$ for $L > 4$.

Fig. 4 Localization length ξ_l versus $|\mu - \mu_c|$ for the 2D Hubbard model. The solid lines correspond to two values of the correlation length exponent: $\nu = 1/4$ and $\nu = 1/2$. The solid circles correspond to the QMC data.

Fig. 5 Same as Fig. 4 but for the 1D Hubbard model.




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5